

## $\mathcal{F}_\phi$ and $\mathcal{F}_g$ Classes

Lino F. Reséndis O.<sup>1</sup> and Luis M. Tovar S.<sup>2</sup>

**Abstract.** In this paper we give a generalization of the Zhao  $F(p, q, s)$ -spaces by using operators instead of functions. In this way we unify and simplify several important results about the classic spaces  $D_p$ ,  $\mathcal{Q}_p$ ,  $B^\alpha$ , etc.

**Mathematics Subject Classification (2000).** 30C45.

**Keywords.**  $\mathcal{Q}_p$  spaces,  $\mathcal{Q}_p^*$  hyperbolic classes,  $\mathcal{Q}_p^\#$  normal classes, Bloch spaces,  $F(p, q, s)$  spaces.

Lino F. Reséndis O.<sup>1</sup>  
Universidad Autónoma Metropolitana, Unidad Azcapotzalco,  
C.B.I. Apartado Postal 16-306 C.P. 02200 México 16,  
D.F. Area de Análisis Matemático y sus Aplicaciones.  
e-mail: lfro@correo.azc.uam.mx

Luis M. Tovar S.<sup>2</sup>  
Escuela Superior de Física y Matemáticas del IPN  
Edif. 9, Unidad ALM, Zacatenco del IPN., C.P. 07300, D.F., México.  
e-mail: tovar@esfm.ipn.mx

Received: April 28, 2008

Accepted: June 10, 2008

---

<sup>1</sup>Partially supported by CONACYT.

<sup>2</sup>Partially supported by CONACYT and COFAA-IPN.